

properties of the two-dimensional Fourier transform

8. Convolution

The *DFT* of convolution of two functions is equal to the product of the DFT of these two functions, i.e.


$$\mathfrak{F} \{ f_1(x, y) * f_2(x, y) \} = F_1(u, v) \cdot F_2(u, v)$$

The convolution of two functions $f(x)$ and $g(x)$ denoted by $f(x)*g(x)$ is defined by the integral:

$$f(x) * g(x) = \int_{-\infty}^{\infty} f(\alpha) g(x - \alpha) d\alpha$$

Where α is a dummy variable of integration. The convolution of two functions $f(x)$ and $g(x)$ before carrying out the integration it is necessary to form the function $g(x - \alpha)$. It is noted that this operation is simply one of folding $g(\alpha)$ about the origin to give $g(-\alpha)$ and then displacing this function by x . Then for any given value of x , we multiply $f(\alpha)$ by the corresponding $g(x - \alpha)$ and integrate the product from $-\infty$ to ∞ .

Example : convolved the following functions:



Solution:


$$f(x) = \begin{cases} 1 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad g(x) = \begin{cases} 1/2 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$f(x) * g(x) = \int_{-\infty}^{\infty} f(\alpha) g(x - \alpha) d\alpha$$


to convolved between two functions we must followed the following steps:

- ***graphically***

step1: change the axis



Step2: folding $g(\alpha)$ about origin to give $g(-\alpha)$ and then displacing this function by x to become $g(x-\alpha)$




Step3: Sliding , for any given value of x , we multiply $f(\alpha)$ by the corresponding $g(x-\alpha)$ and integrate the product from $-\infty$ to ∞ .

(1) when $x < 0$ $f(x) * g(x) = 0$


(2) when $0 \leq x \leq 1$

$$\int_0^x f(\alpha)g(x-\alpha)d\alpha$$

$$= \int_0^x 1 \cdot \frac{1}{2} d\alpha = \frac{1}{2} \alpha \Big|_0^x = \frac{1}{2} x$$




(3) this interval satisfied by step (2) and (4)



(4) when $1 \geq x-1 \geq 0$
 $2 \geq x \geq 1$

$$\int_{x-1}^1 \frac{1}{2} \cdot 1 d\alpha$$


$$= \frac{1}{2} \alpha \Big|_{x-1}^1 = \frac{1}{2} - \frac{1}{2}(x-1) = 1 - \frac{1}{2}x$$



(5) when $x-1 > 1 \Rightarrow x > 2$

$$f(x) * g(x) = 0$$

$$f(x) * g(x) = \begin{cases} 1/2x & 0 \leq x \leq 1 \\ 1 - \frac{1}{2}x & 1 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$



- **mathematically**

$$f(x) * g(x) = \int_{-\infty}^{\infty} f(\alpha)g(x-\alpha)d\alpha$$

$$0 \leq \alpha \leq 1 \text{ for } f(\alpha)$$

$$0 \leq \alpha \leq 1 \text{ for } g(\alpha)$$

$$\begin{matrix} 0 \leq \alpha \leq 1 \\ x-1 \leq \alpha \leq x \end{matrix}$$

$$\int_0^x f(\alpha)g(x-\alpha)d\alpha = \int_0^x 1 \cdot \frac{1}{2}d\alpha = \frac{1}{2}\alpha \Big|_0^x = \frac{1}{2}x$$

$$\int_{x-1}^1 \frac{1}{2} \cdot 1 d\alpha = \frac{1}{2}\alpha \Big|_{x-1}^1 = \frac{1}{2} - \frac{1}{2}(x-1) = 1 - \frac{1}{2}x$$

$$f(x) * g(x) = \begin{cases} 1/2x & 0 \leq x \leq 1 \\ 1 - \frac{1}{2}x & 1 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

