


Chapter Three

Numerical Solution of Ordinary Differential Equations



Theorem (Existence and uniqueness)

Assume that $f(x, y)$ is continuous function in a region $R = \{(x, y) : t_0 \leq t \leq b, c \leq y \leq d\}$. If f satisfies a Lipschitz condition on R in the variable y and $(x_0, y_0) \in R$, then the initial value problem $y' = f(x, y)$ with $y(x_0) = y_0$ has a unique solution $y = y(x)$ on some subinterval $x_0 \leq x \leq x_0 + \delta$.

The Lipschitz condition is

$$|f(x, y_1) - f(x, y_2)| \leq K |y_1 - y_2|, \text{ all } (x, y_1), (x, y_2) \text{ in } R \text{ and}$$
$$|\partial f(x, y) / \partial y| \leq k$$

The general form of differential equation of order (n) is

$$y^{(n)} = f(x, y, y', y'', \dots, y^{(n-1)})$$

In this chapter we solve the ordinary differential equation of order one (Initial value problem) which has the form:

$$y' = f(x, y), \quad y(x_0) = y_0 \tag{1}$$

Let $[a, b]$ be the interval over which we want to find the solution to the well-posed I.V.P. $y' = f(x, y)$ with $y(x_0) = y_0$. In actuality, we will not find a differentiable function that satisfies the I.V.P. Instead, a set of points

$\{(x_k, y_k)\}$ is generated, and the points are used for an approximation

(i.e., $y(x_k) \approx y_k$). How can we proceed to

First we subdivide the interval $[a, b]$ into N equal subintervals and select the mesh points $x_k = a + kh$ for $k = 0, 1, \dots, n$ where $h = (b - a)/n$

The value h is called the step size. We now proceed to solve approximately

Equation (1) over $[x_0, x_n]$ with $y(x_0) = y_0$.

Numerical Methods:

1- Euler's Method

Assume that $y(x)$, $y'(x)$, and $y''(x)$ are continuous and use Taylor's theorem to expand $y(x)$ about $x = x_0$. For each value x there exists a value c_1 that lies between x_0 and x so that

$$y(x) = y(x_0) + y'(x_0)(x - x_0) + y''(c_1)(x - x_0)^2 \quad (2)$$

When $y'(x_0) = f(x_0, y(x_0))$ and $h = x_1 - x_0$ are substituted in eq.(2) the result is an expression for $y(x_1)$:

$$y(x_1) = y(x_0) + h f(x_0, y(x_0)) + y''(c_1) \frac{h^2}{2}$$

i.e.

$$y_1 = y_0 + h f(x_0, y_0)$$

and the local truncation error of Euler method is $O(h^2)$ (L.T.E. = $y''(c_1) \frac{h^2}{2}$)

The process is repeated and generates a sequence of points that approximates the solution curve $y = y(t)$. The general step for Euler's method is

$$\begin{aligned}x_{k+1} &= x_k + h \\y_{k+1} &= y_k + h f(x_k, y_k) \quad \text{for } k = 0, 1, \dots, n-1\end{aligned} \quad (3)$$

Example (1): Use Euler's method to solve approximately the initial value problem

$$\frac{dy(x)}{dx} = x - y, \quad y(0) = 1$$

At the values $x=0.1, 0.2, 0.3, 0.4$

Solution:

$$f(x,y) = x - y, \quad x_0 = 0, \quad y_0 = 1, \quad h = 0.1$$

$$x_{k+1} = x_k + h$$

$$y_{k+1} = y_k + h f(x_k, y_k) \quad \text{for } k = 0, 1, 2, 3$$

$$\begin{aligned}k=0 \longrightarrow y(x_1) &= y(0.1) = y_1 = y_0 + h f(x_0, y_0) \\ &= 1 + (0.1)(x_0 - y_0) = 1 + (0.1)(0 - 1) = 0.9\end{aligned}$$

$$\begin{aligned}k=1 \longrightarrow y(x_2) &= y(0.2) = y_2 = y_1 + h f(x_1, y_1) \\ &= 0.9 + (0.1)(x_1 - y_1) = 0.9 + (0.1)(0.1 - 0.9) = 0.82\end{aligned}$$

$$\begin{aligned}k=2 \longrightarrow y(x_3) &= y(0.3) = y_3 = y_2 + h f(x_2, y_2) \\ &= 0.82 + (0.1)(x_2 - y_2) = 0.82 + (0.1)(0.2 - 0.82) = 0.758\end{aligned}$$

$$\begin{aligned}k=3 \longrightarrow y(x_4) &= y(0.4) = y_4 = y_3 + h f(x_3, y_3) \\ &= 0.758 + (0.1)(x_3 - y_3) = 0.758 + (0.1)(0.3 - 0.758) = 0.7122\end{aligned}$$

The exact solution is $y = 2e^{-x} + x - 1$

| x_k | Euler's approximation $y(x_k)$ | Exact solution | Error |
|-------|-----------------------------------|----------------|----------|
| 0 | 1 | 1 | 0 |
| 0.1 | 0.9 | 0.909675 | 0.009675 |
| 0.2 | 0.82 | 0.837462 | 0.017462 |
| 0.3 | 0.758 | 0.781636 | 0.023636 |
| 0.4 | 0.7122 | 0.74064 | 0.02844 |

Example (2): Use Euler's method to approximate the solution for the initial value problem

$$y' = y - x^2 + 1, \quad y(0) = 0.5, \quad 0 \leq x \leq 1$$

with $n=5$.

Solution:

$$f(x, y) = y - x^2 + 1, \quad x_0=0, \quad y_0=0.5, \quad n=5$$

$$h = \frac{b - a}{n} = \frac{1 - 0}{5} = \frac{1}{5}$$

$$x_{k+1} = x_k + h = x_0 + ih = \frac{1}{5}i$$

$$x_1 = \frac{1}{5}, \quad x_2 = \frac{2}{5}, \quad x_3 = \frac{3}{5}, \quad x_4 = \frac{4}{5}, \quad x_5 = 1$$

$$y_{k+1} = y_k + h f(x_k, y_k) \quad \text{for } k = 0, 1, 2, 3, 4$$

$$k=0 \longrightarrow y(x_1) = y\left(\frac{1}{5}\right) = y_1 = y_0 + h f(x_0, y_0)$$

$$= 0.5 + \left(\frac{1}{5}\right)(y_0 - x_0^2 + 1) = 0.5 + \left(\frac{1}{5}\right)(0.5 - 0 + 1) = 0.8$$

$$k=1 \longrightarrow y(x_2)=y\left(\frac{2}{5}\right)=y_2= y_1 + h f(x_1, y_1) \\ =1.152$$

$$k=2 \longrightarrow y(x_3)=y\left(\frac{3}{5}\right)=y_3= y_2 + h f(x_2, y_2) \\ =01.5504$$

$$k=3 \longrightarrow y(x_4)=y\left(\frac{4}{5}\right)=y_4= y_3 + h f(x_3, y_3) \\ =1.9884$$

$$k=4 \longrightarrow y(x_5)=y(1)=y_5= y_4 + h f(x_4, y_4) \\ =2.4587$$

Dr. E. S. Al-Rawi