

## Shifting and Scaling Graphs

In this section we look at the main ways functions are combined or transformed to form new functions.

### Sums, Differences, Products, and Quotients

Like numbers, functions can be added, subtracted, multiplied, and divided (except where the denominator is zero) to produce new functions. If  $f$  and  $g$  are functions, then for every  $x$  that belongs to the domains of both  $f$  and  $g$  (that is, for  $x \in D(f) \cap D(g)$ ), we define functions  $f + g$ ,  $f - g$ , and  $fg$  by the formulas

$$(f + g)(x) = f(x) + g(x).$$

$$(f - g)(x) = f(x) - g(x).$$

$$(fg)(x) = f(x)g(x).$$

Notice that the  $+$  sign on the left-hand side of the first equation represents the operation of addition of *functions*, whereas the  $+$  on the right-hand side of the equation means addition of the real numbers  $f(x)$  and  $g(x)$ .

At any point of  $D(f) \cap D(g)$  at which  $g(x) \neq 0$ , we can also define the function  $f/g$  by the formula

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad (\text{where } g(x) \neq 0).$$

Functions can also be multiplied by constants: If  $c$  is a real number, then the function  $cf$  is defined for all  $x$  in the domain of  $f$  by

$$(cf)(x) = cf(x).$$

**EXAMPLE 1** The functions defined by the formulas

$$f(x) = \sqrt{x} \quad \text{and} \quad g(x) = \sqrt{1-x}$$


have domains  $D(f) = [0, \infty)$  and  $D(g) = (-\infty, 1]$ . The points common to these domains are the points

$$[0, \infty) \cap (-\infty, 1] = [0, 1].$$


The following table summarizes the formulas and domains for the various algebraic combinations of the two functions. We also write  $f \cdot g$  for the product function  $fg$ .

Function	Formula	Domain
$f + g$	$(f + g)(x) = \sqrt{x} + \sqrt{1-x}$	$[0, 1] = D(f) \cap D(g)$
$f - g$	$(f - g)(x) = \sqrt{x} - \sqrt{1-x}$	$[0, 1]$
$g - f$	$(g - f)(x) = \sqrt{1-x} - \sqrt{x}$	$[0, 1]$
$f \cdot g$	$(f \cdot g)(x) = f(x)g(x) = \sqrt{x(1-x)}$	$[0, 1]$
$f/g$	$\frac{f}{g}(x) = \frac{f(x)}{g(x)} = \sqrt{\frac{x}{1-x}}$	$[0, 1)$ ( $x = 1$ excluded)
$g/f$	$\frac{g}{f}(x) = \frac{g(x)}{f(x)} = \sqrt{\frac{1-x}{x}}$	$(0, 1]$ ( $x = 0$ excluded)

The graph of the function  $f + g$  is obtained from the graphs of  $f$  and  $g$  by adding the corresponding  $y$ -coordinates  $f(x)$  and  $g(x)$  at each point  $x \in D(f) \cap D(g)$ , as in Figure 1.25. The graphs of  $f + g$  and  $f \cdot g$  from Example 1 are shown in Figure 1.26.



**FIGURE 1.25** Graphical addition of two functions.



**FIGURE 1.26** The domain of the function  $f + g$  is the intersection of the domains of  $f$  and  $g$ , the interval  $[0, 1]$  on the  $x$ -axis where these domains overlap. This interval is also the domain of the function  $f \cdot g$  (Example 1).

## Composite Functions


Composition is another method for combining functions.

**DEFINITION** If  $f$  and  $g$  are functions, the **composite** function  $f \circ g$  (“ $f$  composed with  $g$ ”) is defined by


$$(f \circ g)(x) = f(g(x)).$$

The domain of  $f \circ g$  consists of the numbers  $x$  in the domain of  $g$  for which  $g(x)$  lies in the domain of  $f$ .

The definition implies that  $f \circ g$  can be formed when the range of  $g$  lies in the domain of  $f$ . To find  $(f \circ g)(x)$ , *first* find  $g(x)$  and *second* find  $f(g(x))$ . Figure 1.27 pictures  $f \circ g$  as a machine diagram and Figure 1.28 shows the composite as an arrow diagram.



**FIGURE 1.27** Two functions can be composed at  $x$  whenever the value of one function at  $x$  lies in the domain of the other. The composite is denoted by  $f \circ g$ .



**FIGURE 1.28** Arrow diagram for  $f \circ g$ .

To evaluate the composite function  $g \circ f$  (when defined), we find  $f(x)$  first and then  $g(f(x))$ . The domain of  $g \circ f$  is the set of numbers  $x$  in the domain of  $f$  such that  $f(x)$  lies in the domain of  $g$ .

The functions  $f \circ g$  and  $g \circ f$  are usually quite different.

**EXAMPLE 2** If  $f(x) = \sqrt{x}$  and  $g(x) = x + 1$ , find

- (a)  $(f \circ g)(x)$       (b)  $(g \circ f)(x)$       (c)  $(f \circ f)(x)$       (d)  $(g \circ g)(x)$ .

**Solution**

Composite	Domain
(a) $(f \circ g)(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x + 1}$	$[-1, \infty)$
(b) $(g \circ f)(x) = g(f(x)) = f(x) + 1 = \sqrt{x} + 1$	$[0, \infty)$
(c) $(f \circ f)(x) = f(f(x)) = \sqrt{f(x)} = \sqrt{\sqrt{x}} = x^{1/4}$	$[0, \infty)$
(d) $(g \circ g)(x) = g(g(x)) = g(x) + 1 = (x + 1) + 1 = x + 2$	$(-\infty, \infty)$

To see why the domain of  $f \circ g$  is  $[-1, \infty)$ , notice that  $g(x) = x + 1$  is defined for all real  $x$  but belongs to the domain of  $f$  only if  $x + 1 \geq 0$ , that is to say, when  $x \geq -1$ . ■

Notice that if  $f(x) = x^2$  and  $g(x) = \sqrt{x}$ , then  $(f \circ g)(x) = (\sqrt{x})^2 = x$ . However, the domain of  $f \circ g$  is  $[0, \infty)$ , not  $(-\infty, \infty)$ , since  $\sqrt{x}$  requires  $x \geq 0$ .

## Shifting a Graph of a Function

A common way to obtain a new function from an existing one is by adding a constant to each output of the existing function, or to its input variable. The graph of the new function is the graph of the original function shifted vertically or horizontally, as follows.


### Shift Formulas

#### Vertical Shifts

$y = f(x) + k$       Shifts the graph of  $f$  *up*  $k$  units if  $k > 0$   
                                  Shifts it *down*  $|k|$  units if  $k < 0$

#### Horizontal Shifts

$y = f(x + h)$       Shifts the graph of  $f$  *left*  $h$  units if  $h > 0$   
                                  Shifts it *right*  $|h|$  units if  $h < 0$




**FIGURE 1.29** To shift the graph of  $f(x) = x^2$  up (or down), we add positive (or negative) constants to the formula for  $f$  (Examples 3a and b).

## EXAMPLE 3


- (a) Adding 1 to the right-hand side of the formula  $y = x^2$  to get  $y = x^2 + 1$  shifts the graph up 1 unit (Figure 1.29).  
 (b) Adding  $-2$  to the right-hand side of the formula  $y = x^2$  to get  $y = x^2 - 2$  shifts the graph down 2 units (Figure 1.29).  
 (c) Adding 3 to  $x$  in  $y = x^2$  to get  $y = (x + 3)^2$  shifts the graph 3 units to the left (Figure 1.30).  
 (d) Adding  $-2$  to  $x$  in  $y = |x|$ , and then adding  $-1$  to the result, gives  $y = |x - 2| - 1$  and shifts the graph 2 units to the right and 1 unit down (Figure 1.31). ■

## Scaling and Reflecting a Graph of a Function

To scale the graph of a function  $y = f(x)$  is to stretch or compress it, vertically or horizontally. This is accomplished by multiplying the function  $f$ , or the independent variable  $x$ , by an appropriate constant  $c$ . Reflections across the coordinate axes are special cases where  $c = -1$ .



**FIGURE 1.30** To shift the graph of  $y = x^2$  to the left, we add a positive constant to  $x$  (Example 3c). To shift the graph to the right, we add a negative constant to  $x$ .



**FIGURE 1.31** Shifting the graph of  $y = |x|$  2 units to the right and 1 unit down (Example 3d).

### Vertical and Horizontal Scaling and Reflecting Formulas

**For  $c > 1$ , the graph is scaled:**

$y = cf(x)$  Stretches the graph of  $f$  vertically by a factor of  $c$ .

$y = \frac{1}{c}f(x)$  Compresses the graph of  $f$  vertically by a factor of  $c$ .

$y = f(cx)$  Compresses the graph of  $f$  horizontally by a factor of  $c$ .

$y = f(x/c)$  Stretches the graph of  $f$  horizontally by a factor of  $c$ .


**For  $c = -1$ , the graph is reflected:**

$y = -f(x)$  Reflects the graph of  $f$  across the  $x$ -axis.


$y = f(-x)$  Reflects the graph of  $f$  across the  $y$ -axis.

**EXAMPLE 4** Here we scale and reflect the graph of  $y = \sqrt{x}$ .


- (a) **Vertical:** Multiplying the right-hand side of  $y = \sqrt{x}$  by 3 to get  $y = 3\sqrt{x}$  stretches the graph vertically by a factor of 3, whereas multiplying by  $1/3$  compresses the graph by a factor of 3 (Figure 1.32).
- (b) **Horizontal:** The graph of  $y = \sqrt{3x}$  is a horizontal compression of the graph of  $y = \sqrt{x}$  by a factor of 3, and  $y = \sqrt{x/3}$  is a horizontal stretching by a factor of 3 (Figure 1.33). Note that  $y = \sqrt{3x} = \sqrt{3}\sqrt{x}$  so a horizontal compression *may* correspond to a vertical stretching by a different scaling factor. Likewise, a horizontal stretching may correspond to a vertical compression by a different scaling factor.
- (c) **Reflection:** The graph of  $y = -\sqrt{x}$  is a reflection of  $y = \sqrt{x}$  across the  $x$ -axis, and  $y = \sqrt{-x}$  is a reflection across the  $y$ -axis (Figure 1.34). ■



**FIGURE 1.32** Vertically stretching and compressing the graph  $y = \sqrt{x}$  by a factor of 3 (Example 4a).




**FIGURE 1.33** Horizontally stretching and compressing the graph  $y = \sqrt{x}$  by a factor of 3 (Example 4b).



**FIGURE 1.34** Reflections of the graph  $y = \sqrt{x}$  across the coordinate axes (Example 4c).

**EXAMPLE 5** Given the function  $f(x) = x^4 - 4x^3 + 10$  (Figure 1.35a), find formulas to

- (a) compress the graph horizontally by a factor of 2 followed by a reflection across the  $y$ -axis (Figure 1.35b).
- (b) compress the graph vertically by a factor of 2 followed by a reflection across the  $x$ -axis (Figure 1.35c).



**FIGURE 1.35** (a) The original graph of  $f$ . (b) The horizontal compression of  $y = f(x)$  in part (a) by a factor of 2, followed by a reflection across the  $y$ -axis. (c) The vertical compression of  $y = f(x)$  in part (a) by a factor of 2, followed by a reflection across the  $x$ -axis (Example 5).

**Solution**

- (a) We multiply  $x$  by 2 to get the horizontal compression, and by  $-1$  to give reflection across the  $y$ -axis. The formula is obtained by substituting  $-2x$  for  $x$  in the right-hand side of the equation for  $f$ :

$$\begin{aligned} y &= f(-2x) = (-2x)^4 - 4(-2x)^3 + 10 \\ &= 16x^4 + 32x^3 + 10. \end{aligned}$$

- (b) The formula is

$$y = -\frac{1}{2}f(x) = -\frac{1}{2}x^4 + 2x^3 - 5. \quad \blacksquare$$


**Ellipses**

Although they are not the graphs of functions, circles can be stretched horizontally or vertically in the same way as the graphs of functions. The standard equation for a circle of radius  $r$  centered at the origin is


$$x^2 + y^2 = r^2.$$

Substituting  $cx$  for  $x$  in the standard equation for a circle (Figure 1.36a) gives

$$c^2x^2 + y^2 = r^2. \quad (1)$$



**FIGURE 1.36** Horizontal stretching or compression of a circle produces graphs of ellipses.



**FIGURE 1.37** Graph of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ,  $a > b$ , where the major axis is horizontal.

If  $0 < c < 1$ , the graph of Equation (1) horizontally stretches the circle; if  $c > 1$  the circle is compressed horizontally. In either case, the graph of Equation (1) is an ellipse (Figure 1.36). Notice in Figure 1.36 that the  $y$ -intercepts of all three graphs are always  $-r$  and  $r$ . In Figure 1.36b, the line segment joining the points  $(\pm r/c, 0)$  is called the **major axis** of the ellipse; the **minor axis** is the line segment joining  $(0, \pm r)$ . The axes of the ellipse are reversed in Figure 1.36c: The major axis is the line segment joining the points  $(0, \pm r)$ , and the minor axis is the line segment joining the points  $(\pm r/c, 0)$ . In both cases, the major axis is the longer line segment.

If we divide both sides of Equation (1) by  $r^2$ , we obtain

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (2)$$

where  $a = r/c$  and  $b = r$ . If  $a > b$ , the major axis is horizontal; if  $a < b$ , the major axis is vertical. The **center** of the ellipse given by Equation (2) is the origin (Figure 1.37).

Substituting  $x - h$  for  $x$ , and  $y - k$  for  $y$ , in Equation (2) results in

$$\frac{(x - h)^2}{a^2} + \frac{(y - k)^2}{b^2} = 1. \quad (3)$$

Equation (3) is the **standard equation of an ellipse** with center at  $(h, k)$ . The geometric definition and properties of ellipses are reviewed in Section 11.6.

## Exercises 1.2

### Algebraic Combinations

In Exercises 1 and 2, find the domains and ranges of  $f$ ,  $g$ ,  $f + g$ , and  $f \cdot g$ .

1.  $f(x) = x$ ,  $g(x) = \sqrt{x - 1}$
2.  $f(x) = \sqrt{x + 1}$ ,  $g(x) = \sqrt{x - 1}$

In Exercises 3 and 4, find the domains and ranges of  $f$ ,  $g$ ,  $f/g$ , and  $g/f$ .

3.  $f(x) = 2$ ,  $g(x) = x^2 + 1$
4.  $f(x) = 1$ ,  $g(x) = 1 + \sqrt{x}$

### Composites of Functions

5. If  $f(x) = x + 5$  and  $g(x) = x^2 - 3$ , find the following.

- |               |              |
|---------------|--------------|
| a. $f(g(0))$  | b. $g(f(0))$ |
| c. $f(g(x))$  | d. $g(f(x))$ |
| e. $f(f(-5))$ | f. $g(g(2))$ |
| g. $f(f(x))$  | h. $g(g(x))$ |

6. If  $f(x) = x - 1$  and  $g(x) = 1/(x + 1)$ , find the following.

- |                |                |
|----------------|----------------|
| a. $f(g(1/2))$ | b. $g(f(1/2))$ |
| c. $f(g(x))$   | d. $g(f(x))$   |
| e. $f(f(2))$   | f. $g(g(2))$   |
| g. $f(f(x))$   | h. $g(g(x))$   |

In Exercises 7–10, write a formula for  $f \circ g \circ h$ .

7.  $f(x) = x + 1$ ,  $g(x) = 3x$ ,  $h(x) = 4 - x$
8.  $f(x) = 3x + 4$ ,  $g(x) = 2x - 1$ ,  $h(x) = x^2$

$$9. f(x) = \sqrt{x + 1}, \quad g(x) = \frac{1}{x + 4}, \quad h(x) = \frac{1}{x}$$

$$10. f(x) = \frac{x + 2}{3 - x}, \quad g(x) = \frac{x^2}{x^2 + 1}, \quad h(x) = \sqrt{2 - x}$$

Let  $f(x) = x - 3$ ,  $g(x) = \sqrt{x}$ ,  $h(x) = x^3$ , and  $j(x) = 2x$ . Express each of the functions in Exercises 11 and 12 as a composite involving one or more of  $f$ ,  $g$ ,  $h$ , and  $j$ .

- |                           |                         |
|---------------------------|-------------------------|
| 11. a. $y = \sqrt{x} - 3$ | b. $y = 2\sqrt{x}$      |
| c. $y = x^{1/4}$          | d. $y = 4x$             |
| e. $y = \sqrt{(x - 3)^3}$ | f. $y = (2x - 6)^3$     |
| 12. a. $y = 2x - 3$       | b. $y = x^{3/2}$        |
| c. $y = x^9$              | d. $y = x - 6$          |
| e. $y = 2\sqrt{x - 3}$    | f. $y = \sqrt{x^3 - 3}$ |

13. Copy and complete the following table.

$g(x)$	$f(x)$	$(f \circ g)(x)$
a. $x - 7$	$\sqrt{x}$	?
b. $x + 2$	$3x$	?
c. ?	$\sqrt{x - 5}$	$\sqrt{x^2 - 5}$
d. $\frac{x}{x - 1}$	$\frac{x}{x - 1}$	?
e. ?	$1 + \frac{1}{x}$	$x$
f. $\frac{1}{x}$	?	$x$

14. Copy and complete the following table.

	$g(x)$	$f(x)$	$(f \circ g)(x)$
a.	$\frac{1}{x-1}$	$ x $	?
b.	?	$\frac{x-1}{x}$	$\frac{x}{x+1}$
c.	?	$\sqrt{x}$	$ x $
d.	$\sqrt{x}$	?	$ x $

15. Evaluate each expression using the given table of values

$x$	-2	-1	0	1	2
$f(x)$	1	0	-2	1	2
$g(x)$	2	1	0	-1	0

- a.  $f(g(-1))$       b.  $g(f(0))$       c.  $f(f(-1))$   
d.  $g(g(2))$       e.  $g(f(-2))$       f.  $f(g(1))$

16. Evaluate each expression using the functions

$$f(x) = 2 - x, \quad g(x) = \begin{cases} -x, & -2 \leq x < 0 \\ x - 1, & 0 \leq x \leq 2. \end{cases}$$

- a.  $f(g(0))$       b.  $g(f(3))$       c.  $g(g(-1))$   
d.  $f(f(2))$       e.  $g(f(0))$       f.  $f(g(1/2))$

In Exercises 17 and 18, (a) write formulas for  $f \circ g$  and  $g \circ f$  and find the (b) domain and (c) range of each.

17.  $f(x) = \sqrt{x+1}$ ,  $g(x) = \frac{1}{x}$


18.  $f(x) = x^2$ ,  $g(x) = 1 - \sqrt{x}$

19. Let  $f(x) = \frac{x}{x-2}$ . Find a function  $y = g(x)$  so that  
 $(f \circ g)(x) = x$ .


20. Let  $f(x) = 2x^3 - 4$ . Find a function  $y = g(x)$  so that  
 $(f \circ g)(x) = x + 2$ .

### Shifting Graphs

21. The accompanying figure shows the graph of  $y = -x^2$  shifted to two new positions. Write equations for the new graphs.




22. The accompanying figure shows the graph of  $y = x^2$  shifted to two new positions. Write equations for the new graphs.




23. Match the equations listed in parts (a)–(d) to the graphs in the accompanying figure.

- a.  $y = (x-1)^2 - 4$       b.  $y = (x-2)^2 + 2$   
c.  $y = (x+2)^2 + 2$       d.  $y = (x+3)^2 - 2$



24. The accompanying figure shows the graph of  $y = -x^2$  shifted to four new positions. Write an equation for each new graph.




Exercises 25–34 tell how many units and in what directions the graphs of the given equations are to be shifted. Give an equation for the shifted graph. Then sketch the original and shifted graphs together, labeling each graph with its equation.

25.  $x^2 + y^2 = 49$  Down 3, left 2
26.  $x^2 + y^2 = 25$  Up 3, left 4
27.  $y = x^3$  Left 1, down 1
28.  $y = x^{2/3}$  Right 1, down 1
29.  $y = \sqrt{x}$  Left 0.81
30.  $y = -\sqrt{x}$  Right 3
31.  $y = 2x - 7$  Up 7
32.  $y = \frac{1}{2}(x + 1) + 5$  Down 5, right 1
33.  $y = 1/x$  Up 1, right 1
34.  $y = 1/x^2$  Left 2, down 1

Graph the functions in Exercises 35–54.


35.  $y = \sqrt{x + 4}$
36.  $y = \sqrt{9 - x}$
37.  $y = |x - 2|$
38.  $y = |1 - x| - 1$
39.  $y = 1 + \sqrt{x - 1}$
40.  $y = 1 - \sqrt{x}$
41.  $y = (x + 1)^{2/3}$
42.  $y = (x - 8)^{2/3}$
43.  $y = 1 - x^{2/3}$
44.  $y + 4 = x^{2/3}$
45.  $y = \sqrt[3]{x - 1} - 1$
46.  $y = (x + 2)^{3/2} + 1$
47.  $y = \frac{1}{x - 2}$
48.  $y = \frac{1}{x} - 2$
49.  $y = \frac{1}{x} + 2$
50.  $y = \frac{1}{x + 2}$
51.  $y = \frac{1}{(x - 1)^2}$
52.  $y = \frac{1}{x^2} - 1$
53.  $y = \frac{1}{x^2} + 1$
54.  $y = \frac{1}{(x + 1)^2}$

55. The accompanying figure shows the graph of a function  $f(x)$  with domain  $[0, 2]$  and range  $[0, 1]$ . Find the domains and ranges of the following functions, and sketch their graphs.



- a.  $f(x) + 2$
- b.  $f(x) - 1$
- c.  $2f(x)$
- d.  $-f(x)$
- e.  $f(x + 2)$
- f.  $f(x - 1)$
- g.  $f(-x)$
- h.  $-f(x + 1) + 1$

56. The accompanying figure shows the graph of a function  $g(t)$  with domain  $[-4, 0]$  and range  $[-3, 0]$ . Find the domains and ranges of the following functions, and sketch their graphs.



- a.  $g(-t)$
- b.  $-g(t)$
- c.  $g(t) + 3$
- d.  $1 - g(t)$
- e.  $g(-t + 2)$
- f.  $g(t - 2)$
- g.  $g(1 - t)$
- h.  $-g(t - 4)$

### Vertical and Horizontal Scaling

Exercises 57–66 tell by what factor and direction the graphs of the given functions are to be stretched or compressed. Give an equation for the stretched or compressed graph.

57.  $y = x^2 - 1$ , stretched vertically by a factor of 3
58.  $y = x^2 - 1$ , compressed horizontally by a factor of 2
59.  $y = 1 + \frac{1}{x^2}$ , compressed vertically by a factor of 2
60.  $y = 1 + \frac{1}{x^2}$ , stretched horizontally by a factor of 3
61.  $y = \sqrt{x + 1}$ , compressed horizontally by a factor of 4
62.  $y = \sqrt{x + 1}$ , stretched vertically by a factor of 3
63.  $y = \sqrt{4 - x^2}$ , stretched horizontally by a factor of 2
64.  $y = \sqrt{4 - x^2}$ , compressed vertically by a factor of 3
65.  $y = 1 - x^3$ , compressed horizontally by a factor of 3
66.  $y = 1 - x^3$ , stretched horizontally by a factor of 2

### Graphing

In Exercises 67–74, graph each function, not by plotting points, but by starting with the graph of one of the standard functions presented in Figures 1.14–1.17 and applying an appropriate transformation.

67.  $y = -\sqrt{2x + 1}$
68.  $y = \sqrt{1 - \frac{x}{2}}$
69.  $y = (x - 1)^3 + 2$
70.  $y = (1 - x)^3 + 2$
71.  $y = \frac{1}{2x} - 1$
72.  $y = \frac{2}{x^2} + 1$
73.  $y = -\sqrt[3]{x}$
74.  $y = (-2x)^{2/3}$
75. Graph the function  $y = |x^2 - 1|$ .
76. Graph the function  $y = \sqrt{|x|}$ .

### Ellipses

Exercises 77–82 give equations of ellipses. Put each equation in standard form and sketch the ellipse.

77.  $9x^2 + 25y^2 = 225$
78.  $16x^2 + 7y^2 = 112$
79.  $3x^2 + (y - 2)^2 = 3$
80.  $(x + 1)^2 + 2y^2 = 4$



81.  $3(x - 1)^2 + 2(y + 2)^2 = 6$
82.  $6\left(x + \frac{3}{2}\right)^2 + 9\left(y - \frac{1}{2}\right)^2 = 54$
83. Write an equation for the ellipse  $(x^2/16) + (y^2/9) = 1$  shifted 4 units to the left and 3 units up. Sketch the ellipse and identify its center and major axis.
84. Write an equation for the ellipse  $(x^2/4) + (y^2/25) = 1$  shifted 3 units to the right and 2 units down. Sketch the ellipse and identify its center and major axis.

Combining Functions

85. Assume that  $f$  is an even function,  $g$  is an odd function, and both  $f$  and  $g$  are defined on the entire real line  $\mathbb{R}$ . Which of the following (where defined) are even? odd?

- a.  $fg$
- b.  $f/g$
- c.  $g/f$
- d.  $f^2 = ff$
- e.  $g^2 = gg$
- f.  $f \circ g$
- g.  $g \circ f$
- h.  $f \circ f$
- i.  $g \circ g$

86. Can a function be both even and odd? Give reasons for your answer.
- T 87. (Continuation of Example 1.) Graph the functions  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{1 - x}$  together with their (a) sum, (b) product, (c) two differences, (d) two quotients.
- T 88. Let  $f(x) = x - 7$  and  $g(x) = x^2$ . Graph  $f$  and  $g$  together with  $f \circ g$  and  $g \circ f$ .

Trigonometric Functions

This section reviews radian measure and the basic trigonometric functions.

Angles

Angles are measured in degrees or radians. The number of **radians** in the central angle  $A'CB'$  within a circle of radius  $r$  is defined as the number of “radius units” contained in the arc  $s$  subtended by that central angle. If we denote this central angle by  $\theta$  when measured in radians, this means that  $\theta = s/r$  (Figure 1.38), or

$s = r\theta$

( $\theta$  in radians).

(1)

If the circle is a unit circle having radius  $r = 1$ , then from Figure 1.38 and Equation (1), we see that the central angle  $\theta$  measured in radians is just the length of the arc that the angle cuts from the unit circle. Since one complete revolution of the unit circle is  $360^\circ$  or  $2\pi$  radians, we have

$\pi \text{ radians} = 180^\circ$

(2)

and

$1 \text{ radian} = \frac{180}{\pi} (\approx 57.3) \text{ degrees}$

or

$1 \text{ degree} = \frac{\pi}{180} (\approx 0.017) \text{ radians}.$

Table 1.2 shows the equivalence between degree and radian measures for some basic angles.

TABLE 1.2 Angles measured in degrees and radians

Degrees	−180	−135	−90	−45	0	30	45	60	90	120	135	150	180	270	360
$\theta$ (radians)	$-\pi$	$-\frac{3\pi}{4}$	$-\frac{\pi}{2}$	$-\frac{\pi}{4}$	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$