

The solution set is the open interval $(-3/7, \infty)$ (Figure 1.1b).

- (c) The inequality $6/(x - 1) \geq 5$ can hold only if $x > 1$, because otherwise $6/(x - 1)$ is undefined or negative. Therefore, $(x - 1)$ is positive and the inequality will be preserved if we multiply both sides by $(x - 1)$, and we have

$$\begin{aligned}\frac{6}{x - 1} &\geq 5 \\ 6 &\geq 5x - 5 && \text{Multiply both sides by } (x - 1). \\ 11 &\geq 5x && \text{Add 5 to both sides.} \\ \frac{11}{5} &\geq x. && \text{Or } x \leq \frac{11}{5}.\end{aligned}$$

The solution set is the half-open interval $(1, 11/5]$ (Figure 1.1c). ■

Absolute Value

The **absolute value** of a number x , denoted by $|x|$, is defined by the formula

$$|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0. \end{cases}$$

EXAMPLE 2 Finding Absolute Values

$$|3| = 3, \quad |0| = 0, \quad |-5| = -(-5) = 5, \quad | -|a|| = |a|$$
■

Geometrically, the absolute value of x is the distance from x to 0 on the real number line. Since distances are always positive or 0, we see that $|x| \geq 0$ for every real number x , and $|x| = 0$ if and only if $x = 0$. Also,

$$|x - y| = \text{the distance between } x \text{ and } y$$

on the real line (Figure 1.2).

Since the symbol \sqrt{a} always denotes the *nonnegative* square root of a , an alternate definition of $|x|$ is

$$|x| = \sqrt{x^2}.$$

It is important to remember that $\sqrt{a^2} = |a|$. Do not write $\sqrt{a^2} = a$ unless you already know that $a \geq 0$.

The absolute value has the following properties. (You are asked to prove these properties in the exercises.)

Absolute Value Properties

1. $|-a| = |a|$ A number and its additive inverse or negative have the same absolute value.
2. $|ab| = |a||b|$ The absolute value of a product is the product of the absolute values.
3. $\left|\frac{a}{b}\right| = \frac{|a|}{|b|}$ The absolute value of a quotient is the quotient of the absolute values.
4. $|a + b| \leq |a| + |b|$ The **triangle inequality**. The absolute value of the sum of two numbers is less than or equal to the sum of their absolute values.




FIGURE 1.2 Absolute values give distances between points on the number line.

Note that $|-a| \neq -|a|$. For example, $|-3| = 3$, whereas $-|3| = -3$. If a and b differ in sign, then $|a + b|$ is less than $|a| + |b|$. In all other cases, $|a + b|$ equals $|a| + |b|$. Absolute value bars in expressions like $|-3 + 5|$ work like parentheses: We do the arithmetic inside *before* taking the absolute value.




FIGURE 1.3 $|x| < a$ means x lies between $-a$ and a .

EXAMPLE 3 Illustrating the Triangle Inequality

$$|-3 + 5| = |2| = 2 < |-3| + |5| = 8$$

$$|3 + 5| = |8| = |3| + |5|$$

$$|-3 - 5| = |-8| = 8 = |-3| + |-5|$$

The inequality $|x| < a$ says that the distance from x to 0 is less than the positive number a . This means that x must lie between $-a$ and a , as we can see from Figure 1.3.

The following statements are all consequences of the definition of absolute value and are often helpful when solving equations or inequalities involving absolute values.

Absolute Values and Intervals

If a is any positive number, then

5. $|x| = a$ if and only if $x = \pm a$
6. $|x| < a$ if and only if $-a < x < a$
7. $|x| > a$ if and only if $x > a$ or $x < -a$
8. $|x| \leq a$ if and only if $-a \leq x \leq a$
9. $|x| \geq a$ if and only if $x \geq a$ or $x \leq -a$

The symbol \Leftrightarrow is often used by mathematicians to denote the “if and only if” logical relationship. It also means “implies and is implied by.”



EXAMPLE 4 Solving an Equation with Absolute Values

Solve the equation $|2x - 3| = 7$.

Solution By Property 5, $2x - 3 = \pm 7$, so there are two possibilities:

$2x - 3 = 7$	$2x - 3 = -7$	<small>Equivalent equations without absolute values</small>
$2x = 10$	$2x = -4$	<small>Solve as usual.</small>
$x = 5$	$x = -2$	

The solutions of $|2x - 3| = 7$ are $x = 5$ and $x = -2$.

EXAMPLE 5 Solving an Inequality Involving Absolute Values

Solve the inequality $\left|5 - \frac{2}{x}\right| < 1$.

Solution We have

$$\begin{aligned}
 \left| 5 - \frac{2}{x} \right| < 1 &\Leftrightarrow -1 < 5 - \frac{2}{x} < 1 && \text{Property 6} \\
 &\Leftrightarrow -6 < -\frac{2}{x} < -4 && \text{Subtract 5.} \\
 &\Leftrightarrow 3 > \frac{1}{x} > 2 && \text{Multiply by } -\frac{1}{2}. \\
 &\Leftrightarrow \frac{1}{3} < x < \frac{1}{2}. && \text{Take reciprocals.}
 \end{aligned}$$

Notice how the various rules for inequalities were used here. Multiplying by a negative number reverses the inequality. So does taking reciprocals in an inequality in which both sides are positive. The original inequality holds if and only if $(1/3) < x < (1/2)$. The solution set is the open interval $(1/3, 1/2)$. ■



EXAMPLE 6 Solve the inequality and show the solution set on the real line:

(a) $|2x - 3| \leq 1$

(b) $|2x - 3| \geq 1$

Solution

(a)

$$\begin{aligned}
 |2x - 3| &\leq 1 \\
 -1 &\leq 2x - 3 \leq 1 && \text{Property 8} \\
 2 &\leq 2x \leq 4 && \text{Add 3.} \\
 1 &\leq x \leq 2 && \text{Divide by 2.}
 \end{aligned}$$

The solution set is the closed interval $[1, 2]$ (Figure 1.4a).

(b)

$$\begin{aligned}
 |2x - 3| &\geq 1 \\
 2x - 3 &\geq 1 \quad \text{or} \quad 2x - 3 \leq -1 && \text{Property 9} \\
 x - \frac{3}{2} &\geq \frac{1}{2} \quad \text{or} \quad x - \frac{3}{2} \leq -\frac{1}{2} && \text{Divide by 2.} \\
 x &\geq 2 \quad \text{or} \quad x \leq 1 && \text{Add } \frac{3}{2}.
 \end{aligned}$$

The solution set is $(-\infty, 1] \cup [2, \infty)$ (Figure 1.4b). ■




FIGURE 1.4 The solution sets (a) $[1, 2]$ and (b) $(-\infty, 1] \cup [2, \infty)$ in Example 6.